Colorectal Cancer

Screening. Prevention. Early Detection.



Start screening at




2nd

most common cause of cancer deaths in the US



Percent of cases diagnosed under the age of 50

Percent of patients with family history of Colorectal Cancer (CRC)²



This means that most patients <u>do not</u> have a family history of CRC.

We are all at risk!

Colonoscopy

Which screening method is best for you? 3

Stool Kit

- Every 10 years (without history of polyps or family history of CRC)
- 75% of CRC will be prevented by removing pre-cancerous polyps during a colonoscopy
- Decreases deaths from CRC by 85%
- Some risk of complications including bleeding, side effects to prep, intestinal injury
- Barriers include time off work, transportation, and inconvenient prep

- Screening performed annually
- If screening done annually, 60% of CRC will be prevented and deaths from CRC will be decreased by 79%
- Unfortunately, many patients who choose stool kits do not get screened every year
- False positives possible
- Low risk of complication
- If positive, we would refer for a colonoscopy
- Best for patients at average risk

References:

- 1. https://www.cancer.org/cancer/types/colon-rectal-cancer/about/key-statistics.html
- 2.https://www.cancer.org/cancer/types/colon-rectal-cancer/causes-risks-prevention/risk-factors.html
- 3. https://www.cancer.org/cancer/types/colon-rectal-cancer/detection-diagnosis-staging/acs-recommendations.html





Prevention & Medical Care To Help You Live Better, Longer.