

MGB GASTROENTEROLOGY

10 MAIN STREET, FLORENCE, MA 01062 (413) 586-8910

FAX (855) 583-3395

• DAVID A. BERKMAN, MD • JOSEPH P. TASSONI, MD • DAVID R. KALMAN, MD •
• MICHAL GANZ, MD • VIKRAM BUDHRAJA, MD •

SUPREP

Please follow all instructions: An incomplete prep can lead to a poor exam. If you experience nausea/vomiting, take a 30 minute break and then slowly resume the prep. Please stay well hydrated before and during prep.

7 DAYS BEFORE THE PROCEDURE:

- Have your prescription filled. Please follow the prep instructions listed below, **NOT** the instructions on the prep box.

5 DAYS BEFORE THE PROCEDURE:

- If you typically have less than 2 bowel movements per week then you must purchase an 8.3oz bottle of MiraLAX.
- Begin taking a dose of MiraLAX twice a day until the day before your procedure.
- **AVOID** raw fruits and vegetables, raw or cooked corn, salads, red Jell-O, beets, granola, whole wheat/multigrain products, beans, popcorn, nuts, and seeds.

THE DAY BEFORE THE PROCEDURE:

- **CLEAR LIQUIDS ONLY** such as water, coffee, tea, broth, sodas, apple juice, Gatorade, Jell-O, popsicles, etc.
- Please avoid red and purple liquids, dairy and soy products, and juices with pulp.
- **START YOUR PREP AT 4:00pm**

STEP 1:

- Pour **ONE (1)** 6 ounce bottle of SUPREP liquid into mixing container. Add drinking water to the 16 ounce line on the container and mix.
- Drink **ALL** the liquid in the container within one (1) hour.

STEP 2:

- You **must** drink **TWO (2)** more 16 ounce containers of water within the **next 1 hour.**

THE MORNING OF YOUR PROCEDURE:

- **DRINK THE SECOND DOSE OF PREP 6 HOURS PRIOR TO YOUR PROCEDURE**

STEP 1:

- Pour **ONE (1)** 6 ounce bottle of SUPREP liquid into mixing container. Add drinking water to the 16 ounce line on the container and mix.
- Drink **ALL** the liquid in the container within one (1) hour.

STEP 2:

- You **must** drink **TWO (2)** more 16 ounce containers of water within the **next 1 hour.**
- You **must** finish drinking the final container of water **AT LEAST FOUR HOURS** before your procedure.

NO GUM, HARD CANDY, BREATH MINTS, ETC. AFTER MIDNIGHT

ABSOLUTELY NOTHING BY MOUTH 4 HOURS BEFORE PROCEDURE

BLOOD THINNER INSTRUCTIONS:

DIABETIC INSTRUCTIONS:

Do not take any Advil or Aleve the morning of your procedure

****ALL PIERCINGS, JEWELRY AND CONTACT LENSES MUST BE REMOVED PRIOR TO PROCEDURE****

YOU MUST HAVE A RIDE HOME. You will not be discharged from the Endoscopy Unit unless you are accompanied by a responsible adult who will either drive you home or accompany you home by taxi, Uber, or bus.

REPORT TO: _____ Cooley Dickinson Hospital, 30 Locust Street, Northampton, North Entrance

You will be receiving a blocked call from Assessment at CDH, please answer when called

_____ Valley Medical Group, 31 Hall Drive, Amherst - Ambulatory Surgical Center, Main Floor

_____ Pioneer Valley Surgicenter, 3550 Main Street, Springfield, Suite 103

DAY: _____

DATE: _____

ARRIVAL TIME: _____